

Calculus Summer Math Packet

Congratulations! You made it to Calculus!

This summer math packet is a review of concepts learned in Algebra and Pre-Calculus classes that are needed when you begin your Calculus course in August. Completion of the packet will assure that all students begin the school year on the same page and with equal opportunity to learn and build upon mathematical concepts that should have been learned in previous courses.

Instructions for completing the packet:

- Please print the packet or use loose leaf paper to complete the packet by hand showing all work. Work must be neat and legible.
- Please use your Pre-Calculus notes or the websites provided to help you if you need reminders on how to complete some practice problems.
- Take notes as you complete your work. You will be given a quiz on this material the first week of school.
- Work on the packet with your friends. Help each other. Every student is responsible for knowing the material in this packet when you return in August. We will review as a team and everyone will be expected to participate.
- Bring your packet to our first class together. It will be collected for a grade. Only packets done with paper and pencil will be accepted.

A FRIENDLY SABRE SUGGESTION -

begin your summer with a SCHEDULE for completing your packet.

Helpful Websites:

http://www.mathtv.com/ http://www.purplemath.com/modules/index.htm https://www.khanacademy.org

Helpful for graphing functions:

https://www.education.ti.com/en/resources/family-of-functions

CALCULUS SUMMER PACKET

DUE: First day of school.

This assignment is to be done at your leisure during the summer. It is designed to help you become comfortable with your graphing calculator as well as give you practice in necessary math skills which are basic knowledge for Calculus. It is important that you gain these skills over the summer so that we can spend more time talking about calculus and less time on how to use the calculator and do basic algebra.

Part 1: Graph. Analyze each.

- 1. Parent Function: $y = x^2$ a. $y = x^2 - 5$ b. $v = x^2 + 3$ c. $y = (x - 10)^2$ d. $y = (x+8)^2$ e. $y = 4x^2$ f. $y = 0.25x^2$ g. $y = -x^2$ h. $y = -(x+3)^2 + 6$ i. $y = (x+4)^2 - 8$ j. $y = -2(x+1)^2 + 4$ 1. $y = -3(x+2)^2 - 2$ k. $y = \frac{1}{2}(x-6)^2 - 6$ 2. Parent Function: $y = \sin(x)$ a. $y = \sin(2x)$ b. $y = 2\sin(x)$ c. $y = \sin(x) - 2$ d. $y = \sin(2x) + 2$ 3. Parent Function: $y = \cos(x)$ a. $y = \cos(3x)$ b. $y = \cos\left(\frac{x}{2}\right)$ c. $y = 2\cos(x) + 2$ d. $y = -2\cos(x) - 1$ 4. Parent Function: $y = x^3$ b. $y = -x^3$ a. $v = x^3 + 2$ c. $v = x^3 - 5$ d. $v = -x^3 + 3$ e. $y = (x - 4)^3$ f. $y = (x-1)^3 - 4$ g. $y = -2(x+2)^3 + 1$ h. $v = x^3 + x$ 5. Parent Function: $y = \sqrt{x}$
 - a. $y = \sqrt{x} 2$ b. $y = \sqrt{-x}$ c. $y = \sqrt{x} + 5$ d. $y = \sqrt{6-x}$

e.
$$y = -\sqrt{x}$$

g. $y = \sqrt{x+2}$
i. $y = -2\sqrt{x}$
f. $y = -\sqrt{-x}$
h. $y = \sqrt{2x-6}$
j. $y = -\sqrt{4-x}$

6. Parent Function: $y = \ln x$

a. $y = \ln(x+3)$ b. $y = \ln(x)+3$ c. $y = \ln(x-2)$ d. $y = \ln(-x)$ e. $y = -\ln x$ f. $y = \ln |x|$

h. $y = -3\ln(x) + 1$

b. $y = -\frac{1}{x}$

d. $y = \frac{2}{(5-x)}$

g. $y = \ln(2x) - 4$

7. Parent Function: $y = e^x$

- a. $y = e^{2x}$ b. $y = e^{x-2}$ c. $y = e^{2-x}$ d. $y = e^{2x} + 3$ e. $y = -e^x$ f. $y = e^{-x}$ g. $y = 2 e^x$ h. $y = e^{0.5x}$
- 8. Parent Function: $y = 2^x$
 - a. $y = 3^x$ b. $y = 0.5^x$ c. $y = 3^{-x}$ d. $y = 4^{x-3}$ e. $y = 2^{x-3} + 2$ f. $y = 2^{-x} + 3$
- 9. Parent Function: $y = \frac{1}{x}$

a.
$$y = \frac{1}{(x-2)}$$

c. $y = \frac{1}{(x+4)}$

10. Parent Function: y = |x|

a.
$$y = |x| + 2$$

b. $y = |x-3|$
c. $y = -2|x|$
d. $y = |x-2| + 3$

Part 2: Determine the domain and zeros (nearest 0.001) of each of the following functions. Use a sign chart to determine the intervals where the function is positive and negative. Use your calculator and table for help.

For example f(x) = xDomain: {all reals} Zeros: x=0Sign chart: $-1 \quad 0 \quad 1$

11. Given:
$$f(x) = x^4 - 3x^3 + 2x^2 - 7x - 11$$

- 12. Given: $f(x) = 0.7x^2 + 3.2x + 1.5$
- 13. Given: $f(x) = x^4 8x^2 + 5$
- 14. Given: $f(x) = x^3 + 3x^2 10x 1$
- 15. Given: $f(x) = 100x^3 203x^2 + 103x 1$
- 16. Given: $f(x) = \frac{(x+1)}{(x+2)}$

Part 3: Linear equation: Write the following equations in point-slope $y - y_1 = m(x - x_1)$ form.

- 17. The line containing point (4,-7) and having slope of 2.
- 18. The line containing the point (-13,5) and parallel to 4x + 2y = -7
- 19. The line containing the point (0,-2) and perpendicular to x 4y = -7
- 20. The line containing the point (2,9) and having a slope of 0.
- 21. The perpendicular bisector of the segment between (-5,3) and (12,3)

Part 4: For questions 25-27: Copy and complete each (use your calculator)

- a. Sketch the graph of f(x).
- b. Sketch the graph of |f(x)|.

- c. Sketch the graph of f(|x|).
- d. Sketch the graph of f(2x).
- e. Sketch the graph of $2 \cdot f(x)$.

22.
$$f(x) = x^2 - 5x - 3$$

 $23. f(x) = 2\sin(3x)$

24.
$$f(x) = -x^3 - 2x^2 + 3x - 4$$

25. Given: $\frac{f(x) = 3x + 2}{g(x) = -4x - 2}$ Find the point of intersection.

26. Given:
$$\frac{f(x) = x^2 - 5x + 2}{g(x) = 3 - 2x}$$
Find the coordinates of any point(s) of intersection.

- 27. Given: $f(x) = x^4 7x^3 + 6x^2 + 8x + 9$
 - a. Find the lowest point on the graph.
 - b. Find the highest and lowest values of f(x) over the interval $-10 \le x \le 10$.
 - c. Find the highest and lowest values of f(x) over the interval $-2 \le x \le 6$.

Part 5: Factor each of the expressions

- 28. $x^2 8x + 12$ 29. $3x^2 + 8x + 5$ 30. $x^2 - 25$
- 31. $4x^2 81$

Part 6: Composition of Functions: Given f(x) = 4x - 1 and g(x) = x + 6, find the following

compositions:


35. g(f(x))

Part 7: Simplify Expressions

39. ln (1) 40. e^{0} 41. $e^{1+\ln(x)}$ 42. e^{l3lnx} 43. $\frac{\sqrt{x}}{x}$ 44. e^{ln3}

Part 8: Piecewise function:

- 45. Sketch y = |x| as a piecewise function
- 46. Sketch a graph of the piecewise function $f(x) = \begin{cases} -x^2, -2 \le x < 1\\ -2, & x = 1\\ 3x + 5, 1 < x \le 3 \end{cases}$
- 47. The function f(x) is graphed below. Find the following.



Summer MC Review Pre Calc – domain – range / graphing / solving / trig / factoring / odd-even

1) Which of the following defines a function *f* for which f(-x) = -f(x)?

A) D)	2)	$f(x) = x^{2}$ $f(x) = \log x$ $\ln(x-2) < 0 \text{ if and only if}$	B) E)	$f(x) = \sin x$ $f(x) = e^{x}$	C)	$f(x) = \cos x$
A) D)		$\begin{array}{l} x < 3 \\ x > 2 \end{array}$	B) E)	$\begin{array}{l} 0 < x < 3 \\ x > 3 \end{array}$	C)	2< <i>x</i> <3
	3)	The domain of the function define	d by $f($.	$x = \ln(x^2 - 4)$ is the set of all real	l numbers	s <i>x</i> such that
A) D)		$ x < 2$ $ x \ge 2$	B) E)	$ x \le 2$ x is a real number	C)	<i>x</i> > 2
	4)	The domain of the function f given	n by $f(x)$	$x) = \frac{1}{1 - x^2} \text{ is } \{x : x > 1\}, \text{ what is}$	the range	$e ext{ of } f ?$
A)		$\{x: -\infty < x < -1\}$	B)	$\{x: -\infty < x < 0\}$	C)	$\{x: -\infty < x < 1\}$
D)		$\{x: -1 < x < \infty\}$	E)	$\{x: 0 < x < \infty\}$		
	5)	The graph of $y^2 = x^2 + 9$ is syn I. The <i>x</i> -axis II. The <i>y</i> -axis III. The origin	nmetric to	o which of the following?		
A) D)		I only I and II only	B) E)	II only I, II and III	C)	III only
	6)	What is the domain of the function	nfgiven	by $f(x) = \frac{\sqrt{x^2 - 4}}{x - 3}$		
A) D)		$ \{x : x \neq 3\} $ $ \{x : x \ge 2andx \neq 3\} $	B) E)	$ \{x : x \le 2 \} $ $ \{x : x \ge 2 and x \neq 3 \} $	C)	$\{x : \mid x \mid \ge 2\}$
	7)	Which of the following does NOT	` have a p	eriod of π ?		
• >		$f(x) = \sin\left(1, x\right)$	D)		C)	$f(x) = x^2$

A) $f(x) = \sin\left(\frac{1}{2}x\right)$ B) $f(x) = |\sin x|$ C) $f(x) = \sin^2 x$ D) $f(x) = \tan x$ E) $f(x) = \tan^2 x$

- 8) The graph of which of the following equations has y = 1 as an asymptote?
- A) $y = \ln x$ D) $y = \frac{x^2}{x-1}$ B) $y = \sin x$ E) $y = e^{-x}$
 - 9) The graph of which of the following equations has y = 1 as an asymptote?
- A) $y = \ln x$ D) $y = \frac{x^2}{x-1}$ B) $y = \sin x$ E) $y = e^{-x}$
 - 10) Let f and g be odd functions. If p, r, and s are nonzero functions defined as follows, which must be odd? I. p(x) = f(g(x))
 - II. r(x) = f(x) + g(x)III. s(x) = f(x)g(x)
- A)I onlyB)II onlyC)I and II onlyD)II and III onlyE)I, II and IIIII only

11) The fundamental period of $2\cos(3x)$ is

- A) $\frac{2\pi}{3}$ B) 2π
- D) 2 E) 3

 6π

C)

Name _____

0.1: Monomial Factors

Factor as indicated

1.
$$3x^4 + 4x^3 - x^2 = x^2(\underline{})$$

2. $2\sqrt{x} + 6x^{\frac{3}{2}} = 2\sqrt{x}(\underline{})$

3.
$$e^{-x} - xe^{-x} + 2x^2e^{-x} = e^{-x}(__)$$

4. $x^{-1} - 2 + x = x^{-1}(__)$

5.
$$\frac{x}{2} - 6x^2 = \frac{x}{2}$$
 (_____) 6. $\sin^2 x + \tan x = \sin x$ (_____) 7. $\frac{1}{2x^3 + 4x} = \frac{1}{2x}$ (______)

0.2: Factoring Quadratic Expressions

Factor as indicated.

1.
$$x^2 - 3x + 2 = ()()$$
 2. $x^2 - 121 = ()()$

3.
$$x^2 + 5x - 6 = ()()$$
 4. $x^2 + 5x + 6 = ()()$

5.
$$2x^2 + 5x - 3 = ($$
 $)($ $)$ 6. $e^{2x} + 2 + e^{-2x} = ($ $)($ $)$

7.
$$x^4 - 7x^2 + 12 = ()()()$$
 8. $\sin x - \sin^3 x = ()()()$

0.3: Simplify Complex Fractions

Reduce each expression to lowest terms:

1.
$$\frac{3x+9}{6x}$$
 2. $\frac{x^5}{x^{\frac{1}{2}}}$

3.
$$\frac{(x+1)^3(x-2)+3(x+1)^2}{(x+1)^4}$$
 4.
$$\frac{x^{\frac{1}{2}}-x^{\frac{1}{3}}}{x^{\frac{1}{6}}}$$

5.
$$\frac{\sqrt{x-1}+(x-1)^{\frac{3}{2}}}{\sqrt{x-1}}$$
 6. $\frac{1-(\sin x+\cos x)^2}{2\sin x}$

0.4: Quadratic Formula

For each equation, solve for the indicated expression.

1. $x^2 - 4x - 1 = 0$, for x 2. $2x^2 + x - 3 = 0$, for x

3. $\cos^2 x + 3\cos x + 2 = 0$, for $\cos x$, then for x

0.5: Synthetic Division

Use synthetic division to factor as indicated.

1. $x^3 - 4x^2 + 2x + 1 = (x - 1)($) 2. $2x^3 + 5x + 7 = (x + 1)($)

0.6: Special Products

Factor completely (into linear or irreducible quadratic factors):

1.
$$x^3 - 27$$
 2. $x^3 - 3x^2 + 3x - 1$ **3.** $x^3 + 6x^2 + 12x + 8$

4. $x^4 - 25$ (Note: into three factors)

0.7: Factoring by Grouping

Factor completely (into linear or irreducible quadratic factors):

2. $x^3 + 2x^2 + 3x + 6$ 1. $x^3 + 4x^2 - 2x - 8$

*3. $5\cos^2 x - 5\sin^2 x + \sin x + \cos x$ *4. $\cos^2 x + 4\cos x + 4 - \tan^2 x$

0.8: Simplifying

Rewrite each of the following in simplest form:

1.
$$\frac{(x-1)(x+3)-(x+1)^2}{x+1}$$
 2. $\frac{\sqrt{x^2+1}-\frac{1}{\sqrt{x^2+1}}}{x^2+1}$ 3. $\frac{x^2-5x+6}{x^2-4x+4}$

4.
$$\frac{1}{x+1} - \frac{1}{x-1} - \frac{2}{x^2 - 1}$$

0.9: Rationalizing

Remove the sum or difference from the denominator by multiplying the numerator and denominator by the conjugate of the denominator.

1.
$$\frac{1}{1-\cos x}$$
 2. $\frac{x}{1-\sqrt{x^2+1}}$ 3. $\frac{2}{x+\sqrt{x^2+1}}$

Error	Correct form	Comments
$a-(x-b)\neq a-x-b$	a-(x-b)=a-x+b	Change all signs when distribution negative through parentheses.
$(a+b)^2 \neq a^2 + b^2$	$(a + b)^2 = a^2 + 2ab + b^2$	Don't forget middle term when squaring binomials.
$\left(\frac{1}{2}a\right)\left(\frac{1}{2}b\right) \neq \frac{1}{2}ab$	$\left(\frac{1}{2}a\right)\left(\frac{1}{2}b\right) = \frac{1}{4}(ab)$	1/2 occurs twice as a factor.
$\frac{a}{x+b} \neq \frac{a}{x} + \frac{a}{b}$	Leave as $\frac{a}{x+b}$	Don't add denominators when adding fractions.
$\frac{1}{a} + \frac{1}{b} \neq \frac{1}{a+b}$	$\frac{1}{a} + \frac{1}{b} = \frac{a+b}{ab}$	Use definition for adding fractions.
$\frac{x}{a}{b} \neq \frac{bx}{a}$	$\frac{\frac{x}{a}}{\frac{b}{b}} = \left(\frac{x}{a}\right)\left(\frac{1}{b}\right) = \frac{x}{ab}$	Multiply by reciprocal of the denominator.
$\frac{1}{3x} \neq \frac{1}{3}x$	$\frac{1}{3x} = \frac{1}{3} \cdot \frac{1}{x}$	Use definition for multiplying fractions.
$1/x + 2 \neq \frac{1}{x+2}$	$1/x + 2 = \frac{1}{x} + 2$	Be careful when using a slash to denote division.
$(x^2)^3 \neq x^5$	$(x^2)^3 = x^{2 \cdot 3} = x^6$	Multiply exponents when an exponential form is raised to a power.
$2x^3 \neq (2x)^3$	$2x^3 = 2(x^3)$	Exponents have priority over coefficients.
$\frac{1}{x^2 + x^3} \neq x^{-2} + x^{-3}$	Leave as $\frac{1}{x^2 + x^3}$	Don't shift term-by-term from denominator to numerator.
$\sqrt{5x} \neq 5\sqrt{x}$	$\sqrt{5x} = \sqrt{5}\sqrt{x}$	Radicals apply to every factor inside radical.
$\sqrt{x^2 + a^2} \neq x + a$	Leave as $\sqrt{x^2 + a^2}$	Don't apply radicals term-by-term.
$\frac{a+bx}{a} \neq 1+bx$	$\frac{a+bx}{a} = 1 + \frac{b}{a}x$	Cancel common factor, not common terms.
$\frac{a+ax}{a} \neq a+x$	$\frac{a+ax}{a} = 1 + x$	Factor before canceling.

0.10: Algebraic Errors to Avoid

Extra Topic 1: Function Transformation

If $f(x) = x^2 - 1$, describe in words what the following would do to the graph of f(x).


1.
$$f(x)-4$$
 2. $f(x-4)$ **3.** $-f(x+2)$

4.
$$5f(x)+3$$
 5. $f(2x)$ **6.** $|f(x)|$


Here is a graph of y = f(x).


Sketch the following graphs.


7. y = 2f(x)




10. y = f(x+2)









11. y = |f(x)|





12. y = f |x|



Extra Topic 3: Trigonometry Review

Solve the following problems.

If point *P* is on the terminal side of θ , find all 6 trig functions of θ . Draw a picture.

1.
$$P(-2, 4)$$
 2. $P(\sqrt{5}, -2)$

3. If $\cos \theta = \frac{-5}{13}$, θ is quadrant II, find all 6 trig functions of θ . 4. If $\cot \theta = 3$, θ in quadrant III, find all 6 trig functions of θ .

For problems 3-11, solve each equation on the interval
$$[0, 2\pi)$$
.
3. $\sin x = \frac{1}{2}$
4. $\cos^2 x = \cos x$
5. $2\cos x + \sqrt{3} = 0$

6.
$$4\sin^2 x = 1$$

7. $2\sin^2 x + \sin x = 1$
8. $\cos^2 x + 2\cos x = 3$

9. $2\sin x \cos x + \sin x = 0$ 10. $8\cos^2 x - 2\cos x = 1$ 11. $\sin^2 x - \cos^2 x = 0$